

原子爆弾症経過7年後に発症せる骨髓性白血病の一例
第1表

			臓器穿刺所見						末梢血液所見									
			第1回入院時			再入院時			第1回入院時					再入院 T.E.M.前		再入院 T.E.M. 後		
年 月 日			昭 20.9.5	20.10.6	20.10.2 9	27.10 .17	10.25	11.1	20 10.5	" 10.29	" 11.24	21 1.23	22 4.12	27 10.17	" 11.4	" 11.25		
部位			骨髓	骨髓	骨髓	骨髓	脾	肝		末梢血	末梢血	末梢血	末梢血	末梢血	末梢血	末梢血		
種別			有核細胞数	6.00	1.52	13.00	51.00			白血球(万)	0.83	1.54	0.64	0.88	0.94	16.38	14.46	13.70
赤芽球	大	好塩基	1.2	0.3	0.2	0.4	0	0	赤血球(万)	307	360	414		515	260	242	314	
		多染	3.0	0.3	1.6	0	0	0	Hb.(%)	70	92	86	97	99	66	70	66	
		正色	0	0	0.4	0	0	0	F.I.	1.14	1.27	1.03		0.96	1.27	1.40	1.04	
	正	好塩基	0.2	1.0	0.4	0.8	0.8	0										
		多染	8.0	4.3	14.6	1.6	4.4	0.4										
		正色	4.8	5.7	13.6	0.2	0.4	0										
赤分核		0.6	0	0.2	0	0.2	0											
計		17.8	17.0	31.0	3.0	5.6	0.4											
白血球	骨髓芽球		1.2	0.3	2.4	5.2	6.4	2.4							0.8	1.6	3.0	
	好中球	前骨髓	8.6	0	6.2	19.8	18.4	24.6		0					25.4	19.2	19.8	
		骨髓	5.0	0.7	8.6	12.0	9.2	2.6		0					6.4	6.8	6.4	
		後骨髓	5.4	2.3	5.8	13.8	11.8	11.4		0					8.6	15.2	15.2	
		桿核	9.0	12.0	13.2	28.4	25.6	27.8		5.0	0.5	0	6.5	0	34.0	30.0	28.6	
		分節核	10.4	31.7	2.6	8.0	10.2	17.4		51.5	84.0	62.0	59.0	63.0	15.0	20.4	18.0	
	計		38.4	46.7	36.4	82.0	75.2	83.8		56.5	84.5	62.0	65.5	63.0	89.4	91.6	88.0	
	好酸球	前骨髓	0	0	0	0	0.2	0							0	0.8	0.4	
		骨髓	1.8	0	0.8	2.4	2.0	0.2							0.6	0	0	
		後骨髓	0.6	0	0	1.6	0.6	0							0.2	0	0.4	
桿核		1.0	1.0	0.4	1.2	0	0							0.4	0	0		
分節核		0.4	4.0	0.4	2.0	0.4	1.0		5.5				11.5	0.4	0	0.8		
計		3.8	5.0	1.6	7.2	3.2	1.2		5.5	0	0	0	11.5	1.2	0.8	1.6		
好	前骨髓	0	0	0	0	0	0							0				
	骨髓	0	0	0	0	0	0							0.2				

